PROCHECK-NMR

# Ensemble Ramachandran plots
## ensemble_analyzed (20 models)






cv = Circular Variance (low values signify high clustering of the data points). Accessible Buried
Gf = Average G-factor for the residue (the higher the value the more favourable the conformations) based on analysis of high-res. Xstal structures
Data points coloured according to G-factor: Favourable Unfavourable

ensemble_analyzed_06.ps

PROCHECK-NMR

# Ensemble Ramachandran plots
## ensemble_analyzed (20 models)






cv = Circular Variance (low values signify high clustering of the data points). Accessible Buried
Gf = Average G-factor for the residue (the higher the value the more favourable the conformations) based on analysis of high-res. Xstal structures
Data points coloured according to G-factor: Favourable Unfavourable

# Ensemble Ramachandran plots
## ensemble_analyzed (20 models)






cv = Circular Variance (low values signify high clustering of the data points). Accessible Buried
Gf = Average G-factor for the residue (the higher the value the more favourable the conformations) based on analysis of high-res. Xstal structures
Data points coloured according to G-factor: Favourable Unfavourable

# Ensemble Ramachandran plots
## ensemble_analyzed (20 models)






cv = Circular Variance (low values signify high clustering of the data points). Accessible Buried
Gf = Average G-factor for the residue (the higher the value the more favourable the conformations) based on analysis of high-res. Xstal structures
Data points coloured according to G-factor: Favourable Unfavourable

PROCHECK-NMR

# Ensemble Ramachandran plots
## ensemble_analyzed (20 models)






cv = Circular Variance (low values signify high clustering of the data points). Accessible Buried
Gf = Average G-factor for the residue (the higher the value the more favourable the conformations) based on analysis of high-res. Xstal structures
Data points coloured according to G-factor: Favourable Unfavourable