PROCHECK-NMR

Ensemble Ramachandran plots
ensemble_refined (20 models)



cv = Circular Variance (low values signify high clustering of the data points). Accessible Buried
Gf = Average G-factor for the residue (the higher the value the more favourable the conformations) based on analysis of high-res. Xstal structures
Data points coloured according to G-factor: Favourable Unfavourable

ensemble_refined_06.ps

PROCHECK-NMR

Ensemble Ramachandran plots
ensemble_refined (20 models)



cv = Circular Variance (low values signify high clustering of the data points). Accessible Buried
Gf = Average G-factor for the residue (the higher the value the more favourable the conformations) based on analysis of high-res. Xstal structures
Data points coloured according to G-factor: Favourable Unfavourable

ensemble_refined_06.ps

PROCHECK-NMR

Ensemble Ramachandran plots
ensemble_refined (20 models)



cv = Circular Variance (low values signify high clustering of the data points). Accessible Buried
Gf = Average G-factor for the residue (the higher the value the more favourable the conformations) based on analysis of high-res. Xstal structures
Data points coloured according to G-factor: Favourable Unfavourable

Ensemble Ramachandran plots
ensemble_refined (20 models)



cv = Circular Variance (low values signify high clustering of the data points). Accessible Buried
Gf = Average G-factor for the residue (the higher the value the more favourable the conformations) based on analysis of high-res. Xstal structures
Data points coloured according to G-factor: Favourable Unfavourable

Ensemble Ramachandran plots
ensemble_refined (20 models)



cv = Circular Variance (low values signify high clustering of the data points). Accessible Buried
Gf = Average G-factor for the residue (the higher the value the more favourable the conformations) based on analysis of high-res. Xstal structures
Data points coloured according to G-factor: Favourable Unfavourable