



Full wwPDB NMR Structure Validation Report ⓘ

Apr 26, 2016 – 01:47 PM BST

PDB ID : 1C9Q
Title : AVERAGE NMR SOLUTION STRUCTURE OF THE BIR-2 DOMAIN OF XIAP
Authors : Meadows, R.P.; Fesik, S.W.
Deposited on : 1999-08-03

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/NMRValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Mogul : unknown
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : rb-20027457
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : rb-20027457


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR


The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	114402	11133
Ramachandran outliers	111179	9975
Sidechain outliers	111093	9958

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	117	

2 Ensemble composition and analysis ⓘ

This entry contains 1 models. Identification of well-defined residues and clustering analysis are not possible.

3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 1858 atoms, of which 898 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called APOPTOSIS INHIBITOR IAP HOMOLOG.

Mol	Chain	Residues	Atoms						Trace
1	A	117	Total	C	H	N	O	S	0
			1857	603	898	176	176	4	

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	79	ALA	CYS	ENGINEERED	UNP P98170
A	90	GLY	CYS	ENGINEERED	UNP P98170

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms	
2	A	1	Total	Zn
			1	1

4 Residue-property plots [i](#)

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: APOPTOSIS INHIBITOR IAP HOMOLOG

Chain A: 



5 Refinement protocol and experimental data overview ⓘ

The models were refined using the following method: *TORSION ANGLE DYNAMICS, SIMULATED ANNEALING*.

Of the 20 calculated structures, 1 were deposited, based on the following criterion: *NONE*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	structure solution	3.8X
X-PLOR	refinement	3.8X

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality ⓘ

6.1 Standard geometry ⓘ

Bond lengths and bond angles in the following residue types are not validated in this section:
ZN

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts ⓘ

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	959	898	898	35
All	All	960	898	898	35

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 19.

All clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)
1:A:58:PRO:HA	1:A:61:LEU:HD12	0.71	1.61
1:A:112:LEU:HD12	1:A:113:ASN:N	0.69	2.00
1:A:112:LEU:HD11	1:A:114:ILE:HG23	0.69	1.65
1:A:64:ALA:HB1	1:A:94:TRP:HB3	0.68	1.65
1:A:66:LEU:HD23	1:A:76:GLN:C	0.60	2.17
1:A:69:THR:HG21	1:A:76:GLN:OE1	0.59	1.97
1:A:19:ARG:CD	1:A:20:THR:HG22	0.57	2.30
1:A:56:LEU:HD13	1:A:93:ALA:CA	0.56	2.30
1:A:30:ILE:O	1:A:30:ILE:HG23	0.56	2.00
1:A:51:PRO:HG2	1:A:54:ALA:HB2	0.54	1.79
1:A:18:LEU:CD2	1:A:23:VAL:HG23	0.50	2.37
1:A:19:ARG:HD2	1:A:20:THR:HG22	0.49	1.85
1:A:3:HIS:CD2	1:A:4:PHE:CD1	0.49	3.00

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)
1:A:112:LEU:CD1	1:A:114:ILE:HG23	0.49	2.35
1:A:50:TRP:CZ2	1:A:54:ALA:CB	0.49	2.96
1:A:69:THR:HG23	1:A:71:ILE:HG13	0.48	1.85
1:A:6:LEU:HD12	1:A:7:ASP:N	0.48	2.24
1:A:4:PHE:O	1:A:5:ALA:HB3	0.47	2.09
1:A:56:LEU:HD13	1:A:93:ALA:HB2	0.46	1.88
1:A:24:VAL:HG13	1:A:24:VAL:O	0.46	2.11
1:A:18:LEU:HD22	1:A:23:VAL:HG23	0.45	1.87
1:A:94:TRP:HE3	1:A:107:VAL:HG22	0.45	1.72
1:A:54:ALA:O	1:A:55:HIS:CD2	0.44	2.71
1:A:66:LEU:HD22	1:A:75:VAL:CG2	0.43	2.43
1:A:112:LEU:HD12	1:A:113:ASN:H	0.43	1.72
1:A:13:HIS:CD2	1:A:13:HIS:N	0.43	2.85
1:A:110:ARG:O	1:A:111:ASN:CB	0.42	2.67
1:A:3:HIS:NE2	1:A:4:PHE:CD1	0.42	2.87
1:A:56:LEU:HD13	1:A:93:ALA:CB	0.42	2.45
1:A:54:ALA:O	1:A:55:HIS:CG	0.41	2.73
1:A:106:PHE:CE2	1:A:111:ASN:OD1	0.41	2.73
1:A:8:ARG:CZ	1:A:8:ARG:CB	0.41	2.98
1:A:87:TRP:CH2	1:A:93:ALA:O	0.41	2.74
1:A:6:LEU:HD13	1:A:9:PRO:HD2	0.41	1.92
1:A:68:TYR:CE1	1:A:72:GLY:O	0.40	2.75

6.3 Torsion angles ⓘ

6.3.1 Protein backbone ⓘ

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	115/117 (98%)	73 (63%)	31 (27%)	11 (10%)	2	11
All	All	115/117 (98%)	73 (63%)	31 (27%)	11 (10%)	2	11

All 11 Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
1	A	108	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	11	GLU
1	A	113	ASN
1	A	25	ASP
1	A	115	ARG
1	A	111	ASN
1	A	22	GLN
1	A	30	ILE
1	A	55	HIS
1	A	29	THR
1	A	6	LEU

6.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	100/100 (100%)	72 (72%)	28 (28%)	2	20
All	All	100/100 (100%)	72 (72%)	28 (28%)	2	20

All 28 residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type
1	A	85	LYS
1	A	98	ARG
1	A	45	LYS
1	A	48	GLN
1	A	30	ILE
1	A	92	ARG
1	A	115	ARG
1	A	46	SER
1	A	8	ARG
1	A	96	GLU
1	A	13	HIS
1	A	57	THR
1	A	29	THR
1	A	1	ARG
1	A	63	SER

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	91	ASP
1	A	112	LEU
1	A	111	ASN
1	A	52	ASP
1	A	59	ARG
1	A	3	HIS
1	A	33	ARG
1	A	19	ARG
1	A	25	ASP
1	A	22	GLN
1	A	116	SER
1	A	37	MET
1	A	6	LEU

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided