



# Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 01:14 AM GMT

PDB ID : 2C9U  
Title : 1.24 ANGSTROMS RESOLUTION STRUCTURE OF AS-ISOLATED CU-ZN HUMAN SUPEROXIDE DISMUTASE  
Authors : Strange, R.W.; Antonyuk, S.V.; Hough, M.A.; Doucette, P.A.; Valentine S, J.S.; Hasnain, S.  
Deposited on : 2005-12-14  
Resolution : 1.24 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.  
We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>  
with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7 (RC4), CSD as536be (2015)  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20026688  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk26865


# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

## *X-RAY DIFFRACTION*

The reported resolution of this entry is 1.24 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	1229 (1.28-1.20)
Clashscore	102246	1327 (1.28-1.20)
Ramachandran outliers	100387	1274 (1.28-1.20)
Sidechain outliers	100360	1272 (1.28-1.20)
RSRZ outliers	91569	1233 (1.28-1.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	153	<div> <div>16%</div> <div>90%</div> <div>9%</div> </div>
1	F	153	<div> <div>24%</div> <div>87%</div> <div>12%</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	A	1154	-	-	-	X
2	SO4	A	1155	-	-	-	X
2	SO4	F	1154	-	-	X	-
5	ACT	F	1158	-	-	-	X
5	ACT	F	1159	-	-	X	-

## 2 Entry composition [i](#)

There are 6 unique types of molecules in this entry. The entry contains 2691 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called SUPEROXIDE DISMUTASE [CU-ZN].

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	153	Total	C	N	O	S	0	8	0
			1129	690	208	227	4			
1	F	153	Total	C	N	O	S	0	12	0
			1144	699	207	232	6			

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		
2	A	1	Total	O	S	0	0
			5	4	1		
2	F	1	Total	O	S	0	0
			5	4	1		


- Molecule 3 is COPPER (II) ION (three-letter code: CU) (formula: Cu).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Cu 1 1	0	0
3	F	1	Total Cu 1 1	0	0

- Molecule 4 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	2	Total Zn 2 2	0	0
4	F	2	Total Zn 2 2	0	0

- Molecule 5 is ACETATE ION (three-letter code: ACT) (formula: C<sub>2</sub>H<sub>3</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	F	1	Total C O 4 2 2	0	0
5	F	1	Total C O 4 2 2	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	197	Total O 197 197	0	0

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	F	192	Total 192	O 192	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

#### • Molecule 1: SUPEROXIDE DISMUTASE [CU-ZN]



#### • Molecule 1: SUPEROXIDE DISMUTASE [CU-ZN]



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	38.64Å 67.55Å 52.32Å 90.00° 106.48° 90.00°	Depositor
Resolution (Å)	50.19 – 1.24 19.67 – 1.24	Depositor EDS
% Data completeness (in resolution range)	95.8 (50.19-1.24) 93.9 (19.67-1.24)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.12 (at 1.24Å)	Xtriage
Refinement program	REFMAC 5.2.0005	Depositor
R, $R_{free}$	0.139 , 0.176 0.251 , 0.234	Depositor DCC
$R_{free}$ test set	3458 reflections (5.06%)	DCC
Wilson B-factor (Å <sup>2</sup> )	17.6	Xtriage
Anisotropy	0.156	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.27 , 67.0	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.46$ , $\langle L^2 \rangle = 0.29$	Xtriage
Outliers	0 of 68348 reflections	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	2691	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	16.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.25% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, SO4, CU, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.93	2/1189 (0.2%)	0.90	2/1602 (0.1%)
1	F	1.03	6/1208 (0.5%)	0.92	0/1630
All	All	0.98	8/2397 (0.3%)	0.91	2/3232 (0.1%)

All (8) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	F	49	GLU	CG-CD	-7.05	1.41	1.51
1	F	59[A]	SER	CA-CB	6.97	1.63	1.52
1	F	59[B]	SER	CA-CB	6.97	1.63	1.52
1	F	78	GLU	CD-OE1	6.79	1.33	1.25
1	A	59[A]	SER	CA-CB	6.61	1.62	1.52
1	A	59[B]	SER	CA-CB	6.61	1.62	1.52
1	F	78	GLU	CD-OE2	5.63	1.31	1.25
1	F	78	GLU	CG-CD	5.21	1.59	1.51

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	115	ARG	NE-CZ-NH1	6.03	123.31	120.30
1	A	115	ARG	NE-CZ-NH2	-5.58	117.51	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1129	0	1099	15	0
1	F	1144	0	1116	32	1
2	A	10	0	0	1	0
2	F	5	0	0	2	0
3	A	1	0	0	0	0
3	F	1	0	0	0	0
4	A	2	0	0	0	0
4	F	2	0	0	0	0
5	F	8	0	6	9	0
6	A	197	0	0	9	2
6	F	192	0	0	20	3
All	All	2691	0	2221	51	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (51) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:53[B]:ASN:OD1	6:A:2069:HOH:O	1.57	1.17
1:A:49[B]:GLU:OE1	6:A:2061:HOH:O	1.67	1.08
1:A:40:GLU:OE2	6:A:2049:HOH:O	1.71	1.08
1:F:53[B]:ASN:OD1	6:F:2075:HOH:O	1.70	1.07
2:F:1154:SO4:O4	6:F:2186:HOH:O	1.72	1.06
2:A:1154:SO4:O1	6:A:2191:HOH:O	1.75	1.03
1:F:133:GLU:OE1	6:F:2168:HOH:O	1.76	1.01
1:F:13:PRO:HA	6:F:2011:HOH:O	1.59	1.00
1:F:12:GLY:N	6:F:2011:HOH:O	1.95	0.98
1:A:90:ASP:OD1	1:A:92[A]:ASP:OD1	1.80	0.98
1:F:77:GLU:O	6:F:2109:HOH:O	1.80	0.96
2:F:1154:SO4:O1	6:F:2189:HOH:O	1.85	0.93
1:F:10:GLY:O	6:F:2011:HOH:O	1.85	0.92
1:A:57:CYS:HG	1:A:146:CYS:HG	0.89	0.87
1:F:10:GLY:O	6:F:2012:HOH:O	1.96	0.83
1:F:121:GLU:OE1	1:F:122:LYS:NZ	2.14	0.80
1:F:11:ASP:C	6:F:2011:HOH:O	2.18	0.80
1:F:11:ASP:N	5:F:1159:ACT:C	2.46	0.78
1:F:11:ASP:CA	6:F:2011:HOH:O	2.32	0.76
1:F:10:GLY:HA2	5:F:1159:ACT:O	1.84	0.76

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:49:GLU:HG3	6:F:2068:HOH:O	1.84	0.75
1:F:11:ASP:H	5:F:1159:ACT:C	2.00	0.75
1:A:53[B]:ASN:ND2	6:A:2070:HOH:O	2.00	0.71
1:A:52:ASP:OD2	6:A:2064:HOH:O	2.09	0.70
1:A:109:ASP:O	1:A:109:ASP:OD1	2.16	0.64
1:A:136:LYS:NZ	6:A:2176:HOH:O	2.29	0.64
1:F:11:ASP:HA	6:F:2011:HOH:O	1.97	0.62
1:A:36:LYS:HE3	6:A:2012:HOH:O	2.01	0.60
1:A:109:ASP:C	1:A:109:ASP:OD1	2.39	0.60
1:A:90:ASP:CG	1:A:92[A]:ASP:OD1	2.41	0.59
1:F:11:ASP:H	5:F:1159:ACT:CH3	2.19	0.55
1:F:10:GLY:CA	5:F:1159:ACT:O	2.53	0.55
1:F:128:LYS:HD3	6:F:2165:HOH:O	2.07	0.55
1:A:57:CYS:HG	1:A:146:CYS:CB	2.20	0.55
1:F:11:ASP:H	5:F:1159:ACT:H2	1.73	0.54
1:F:30[A]:LYS:HD3	6:F:2040:HOH:O	2.06	0.54
1:F:11:ASP:N	5:F:1159:ACT:O	2.41	0.52
1:F:30[A]:LYS:CD	6:F:2040:HOH:O	2.58	0.52
1:F:121:GLU:CD	1:F:122:LYS:HZ1	2.15	0.50
1:F:30[A]:LYS:HG3	1:F:100:GLU:HG2	1.92	0.50
1:F:118:VAL:HG22	1:F:146[B]:CYS:SG	2.52	0.50
1:F:121:GLU:CD	1:F:122:LYS:NZ	2.66	0.49
1:F:10:GLY:N	6:F:2012:HOH:O	2.27	0.47
1:F:10:GLY:C	5:F:1159:ACT:O	2.53	0.46
1:A:30[A]:LYS:HG2	6:A:2141:HOH:O	2.14	0.46
1:F:30[B]:LYS:HG3	6:F:2135:HOH:O	2.14	0.46
1:F:24:GLU:HG3	1:F:26:ASN:OD1	2.17	0.44
1:F:49:GLU:CG	6:F:2068:HOH:O	2.56	0.44
5:F:1159:ACT:H3	6:F:2175:HOH:O	2.20	0.41
1:A:38:LEU:O	1:A:93:GLY:HA2	2.21	0.41
1:F:10:GLY:HA3	1:F:144:LEU:O	2.21	0.41

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:2110:HOH:O	6:F:2067:HOH:O[2_746]	1.96	0.24
6:A:2058:HOH:O	6:F:2124:HOH:O[2_645]	2.08	0.12
1:F:26:ASN:ND2	6:F:2012:HOH:O[1_655]	2.10	0.10

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	159/153 (104%)	158 (99%)	1 (1%)	0	100	100
1	F	162/153 (106%)	161 (99%)	1 (1%)	0	100	100
All	All	321/306 (105%)	319 (99%)	2 (1%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	124/118 (105%)	121 (98%)	3 (2%)	57	15
1	F	129/118 (109%)	129 (100%)	0	100	100
All	All	253/236 (107%)	250 (99%)	3 (1%)	84	43

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	92[A]	ASP
1	A	92[B]	ASP
1	A	109	ASP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	26	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

Of 11 ligands modelled in this entry, 6 are monoatomic - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
2	SO4	A	1154	4	4,4,4	1.08	0	6,6,6	0.37	0
2	SO4	A	1155	-	4,4,4	0.66	0	6,6,6	0.70	0
2	SO4	F	1154	4	4,4,4	0.89	0	6,6,6	0.47	0
5	ACT	F	1158	-	1,3,3	6.16	1 (100%)	0,3,3	0.00	-
5	ACT	F	1159	1	1,3,3	1.62	0	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	1154	4	-	0/0/0/0	0/0/0/0

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	1155	-	-	0/0/0/0	0/0/0/0
2	SO4	F	1154	4	-	0/0/0/0	0/0/0/0
5	ACT	F	1158	-	-	0/0/0/0	0/0/0/0
5	ACT	F	1159	1	-	0/0/0/0	0/0/0/0

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	F	1158	ACT	CH3-C	6.16	1.57	1.48

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	1154	SO4	1	0
2	F	1154	SO4	2	0
5	F	1159	ACT	9	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	153/153 (100%)	1.21	25 (16%) <b>2</b> <b>3</b>	7, 12, 23, 29	23 (15%)
1	F	153/153 (100%)	1.39	36 (23%) <b>1</b> <b>1</b>	8, 13, 21, 28	26 (16%)
All	All	306/306 (100%)	1.30	61 (19%) <b>1</b> <b>2</b>	7, 12, 21, 29	49 (16%)

All (61) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	F	10	GLY	6.6
1	F	1[A]	ALA	6.6
1	F	132	GLU	6.2
1	A	1	ALA	6.0
1	A	130	GLY	5.9
1	F	11	ASP	5.6
1	F	130	GLY	5.1
1	F	134	SER	4.7
1	F	131	ASN	4.7
1	F	13	PRO	4.1
1	F	26	ASN	4.0
1	F	135	THR	3.7
1	F	84	LEU	3.7
1	A	132	GLU	3.5
1	A	67	LEU	3.3
1	A	14	VAL	3.2
1	F	127	GLY	3.1
1	A	119	VAL	3.0
1	A	84	LEU	3.0
1	A	11	ASP	3.0
1	F	144	LEU	2.9
1	F	94	VAL	2.8
1	A	133	GLU	2.8
1	A	99	ILE	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	F	25	SER	2.6
1	F	137	THR	2.6
1	A	38	LEU	2.6
1	A	5	VAL	2.6
1	F	109	ASP	2.6
1	F	87	VAL	2.5
1	F	126	LEU	2.5
1	A	57	CYS	2.5
1	F	108	GLY	2.5
1	A	126	LEU	2.4
1	A	39	THR	2.4
1	F	117	LEU	2.4
1	F	12	GLY	2.4
1	A	47	VAL	2.4
1	A	131	ASN	2.4
1	A	134	SER	2.3
1	F	32	TRP	2.3
1	A	94	VAL	2.3
1	F	140	ALA	2.3
1	A	135	THR	2.3
1	F	146[A]	CYS	2.3
1	A	137	THR	2.2
1	F	92	ASP	2.2
1	F	96[A]	ASP	2.2
1	F	93	GLY	2.2
1	A	2	THR	2.2
1	A	87	VAL	2.1
1	F	104	ILE	2.1
1	F	107	SER	2.1
1	F	45	PHE	2.1
1	F	142	SER	2.1
1	A	117	LEU	2.1
1	A	104	ILE	2.1
1	F	136	LYS	2.1
1	F	133	GLU	2.0
1	F	47	VAL	2.0
1	F	81	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.



### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors( $\text{\AA}^2$ )	Q<0.9
5	ACT	F	1158	4/4	0.64	0.33	6.65	5,12,18,18	4
2	SO4	A	1155	5/5	0.92	0.21	3.96	21,25,27,28	5
2	SO4	A	1154	5/5	0.90	0.18	2.59	7,9,14,14	5
2	SO4	F	1154	5/5	0.92	0.13	-0.17	14,16,19,20	5
4	ZN	F	1157	1/1	0.98	0.05	-1.19	9,9,9,9	0
4	ZN	A	1158	1/1	0.97	0.07	-1.56	9,9,9,9	0
4	ZN	F	1155	1/1	0.96	0.07	-1.77	11,11,11,11	1
4	ZN	A	1159	1/1	1.00	0.03	-7.46	9,9,9,9	1
3	CU	A	1157	1/1	1.00	0.03	-7.91	10,10,10,10	1
5	ACT	F	1159	4/4	0.57	0.24	-	18,21,23,23	4
3	CU	F	1156	1/1	0.98	0.11	-	9,9,9,9	1

### 6.5 Other polymers [i](#)

There are no such residues in this entry.