



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 04:34 PM GMT

PDB ID : 4FEU
Title : Crystal structure of the aminoglycoside phosphotransferase APH(3')-Ia, with substrate kanamycin and small molecule inhibitor anthracycline SP600125
Authors : Stogios, P.J.; Evdokimova, E.; Wawrzak, Z.; Minasov, G.; Egorova, O.; Di Leo, R.; Shakyia, T.; Spanogiannopoulos, P.; Wright, G.D.; Savchenko, A.; Anderson, W.F.; Center for Structural Genomics of Infectious Diseases (CSGID)
Deposited on : 2012-05-30
Resolution : 2.37 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.37 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	4019 (2.40-2.36)
Clashscore	102246	4595 (2.40-2.36)
Ramachandran outliers	100387	4520 (2.40-2.36)
Sidechain outliers	100360	4522 (2.40-2.36)
RSRZ outliers	91569	4034 (2.40-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	272	<div> <div>2%</div> <div>82% 10% 7%</div> </div>
1	B	272	<div> <div>8%</div> <div>88% 8%</div> </div>
1	C	272	<div> <div>6%</div> <div>92% 5%</div> </div>
1	D	272	<div> <div>14%</div> <div>78% 10% 11%</div> </div>
1	E	272	<div> <div>4%</div> <div>91% 6%</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
1	F	272	<div><div></div><div>3%</div><div>86%</div><div>•</div><div>10%</div></div>

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 13331 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Aminoglycoside 3'-phosphotransferase AphA1-IAB.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	253	Total	C	N	O	S	Se	0	1	0
			2042	1307	348	377	4	6			
1	B	263	Total	C	N	O	S	Se	0	2	0
			2138	1362	373	392	5	6			
1	C	262	Total	C	N	O	S	Se	0	1	0
			2116	1348	364	393	5	6			
1	D	242	Total	C	N	O	S	Se	0	0	0
			1952	1252	334	356	4	6			
1	E	264	Total	C	N	O	S	Se	0	0	0
			2125	1354	368	392	5	6			
1	F	245	Total	C	N	O	S	Se	0	2	0
			2000	1282	342	366	4	6			

There are 6 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	0	GLY	-	EXPRESSION TAG	UNP B0VD92
B	0	GLY	-	EXPRESSION TAG	UNP B0VD92
C	0	GLY	-	EXPRESSION TAG	UNP B0VD92
D	0	GLY	-	EXPRESSION TAG	UNP B0VD92
E	0	GLY	-	EXPRESSION TAG	UNP B0VD92
F	0	GLY	-	EXPRESSION TAG	UNP B0VD92

- Molecule 2 is KANAMYCIN A (three-letter code: KAN) (formula: C₁₈H₃₆N₄O₁₁).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
2	A	1	Total	C	N	O	0	0
			33	18	4	11		
2	B	1	Total	C	N	O	0	0
			33	18	4	11		
2	C	1	Total	C	N	O	0	0
			33	18	4	11		
2	D	1	Total	C	N	O	0	0
			33	18	4	11		
2	E	1	Total	C	N	O	0	0
			33	18	4	11		
2	F	1	Total	C	N	O	0	0
			33	18	4	11		

- Molecule 3 is 2,6-DIHYDROANTHRA/1,9-CD/PYRAZOL-6-ONE (three-letter code: 537) (formula: C₁₄H₈N₂O).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
3	A	1	Total	C	N	O	0	0
			17	14	2	1		
3	C	1	Total	C	N	O	0	0
			17	14	2	1		
3	E	1	Total	C	N	O	0	0
			17	14	2	1		

- Molecule 4 is ACETATE ION (three-letter code: ACT) (formula: $C_2H_3O_2$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	C	1	Total	C	O	0	0
			4	2	2		
4	D	1	Total	C	O	0	0
			4	2	2		
4	F	1	Total	C	O	0	0
			4	2	2		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	129	Total	O	0	0
			129	129		
5	B	135	Total	O	0	3
			138	138		
5	C	129	Total	O	0	1
			130	130		
5	D	45	Total	O	0	0
			45	45		
5	E	126	Total	O	0	3
			129	129		
5	F	111	Total	O	0	3
			114	114		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Aminoglycoside 3'-phosphotransferase AphA1-IAB




- Molecule 1: Aminoglycoside 3'-phosphotransferase AphA1-IAB




- Molecule 1: Aminoglycoside 3'-phosphotransferase AphA1-IAB




- Molecule 1: Aminoglycoside 3'-phosphotransferase AphA1-IAB







- Molecule 1: Aminoglycoside 3'-phosphotransferase AphA1-IAB



- Molecule 1: Aminoglycoside 3'-phosphotransferase AphA1-IAB



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	57.75Å 94.16Å 96.77Å 61.21° 73.11° 87.36°	Depositor
Resolution (Å)	19.97 – 2.37 19.97 – 2.37	Depositor EDS
% Data completeness (in resolution range)	(Not available) (19.97-2.37) 80.9 (19.97-2.37)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.07	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.37 (at 2.38Å)	Xtriage
Refinement program	BUSTER 2.8.0	Depositor
R, R_{free}	0.160 , 0.215 0.164 , 0.218	Depositor DCC
R_{free} test set	3406 reflections (5.32%)	DCC
Wilson B-factor (Å ²)	40.5	Xtriage
Anisotropy	0.333	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.31 , 56.2	EDS
Estimated twinning fraction	0.010 for -h,-k,-h-k+l	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtriage
Outliers	0 of 67481 reflections	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	13331	wwPDB-VP
Average B, all atoms (Å ²)	55.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.81% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 537, KAN, ACT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.50	0/2090	0.68	0/2828
1	B	0.49	0/2186	0.66	0/2952
1	C	0.51	0/2164	0.68	0/2925
1	D	0.48	0/1998	0.63	0/2700
1	E	0.50	0/2173	0.67	0/2937
1	F	0.49	0/2046	0.67	0/2762
All	All	0.49	0/12657	0.67	0/17104

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2042	0	1978	16	0
1	B	2138	0	2076	15	0
1	C	2116	0	2045	6	0
1	D	1952	0	1896	19	0
1	E	2125	0	2062	10	0
1	F	2000	0	1937	6	0
2	A	33	0	36	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	B	33	0	36	0	0
2	C	33	0	36	0	0
2	D	33	0	36	0	0
2	E	33	0	36	0	0
2	F	33	0	36	0	0
3	A	17	0	8	0	0
3	C	17	0	8	0	0
3	E	17	0	8	0	0
4	A	4	0	3	0	0
4	B	4	0	3	0	0
4	C	8	0	6	0	0
4	D	4	0	3	0	0
4	F	4	0	3	0	0
5	A	129	0	0	1	0
5	B	138	0	0	1	0
5	C	130	0	0	0	0
5	D	45	0	0	0	0
5	E	129	0	0	1	0
5	F	114	0	0	1	0
All	All	13331	0	12252	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:258:MSE:SE	1:A:261:LEU:HD23	2.38	0.74
1:E:34:GLY:H	1:E:35:GLN:HG3	1.58	0.68
1:A:234:ASN:HB2	1:A:265:LEU:HD11	1.81	0.62
1:E:256:PRO:HB2	1:E:258:MSE:CE	2.34	0.58
1:B:124:LEU:HD23	1:B:211:LEU:HD21	1.85	0.57
1:C:181:GLU:HA	1:C:184:LYS:HE3	1.87	0.56
1:C:70:VAL:HG11	1:D:70:VAL:HG11	1.87	0.56
1:D:234:ASN:HB2	1:D:265:LEU:HD11	1.89	0.54
1:D:46:LYS:HB3	1:D:49:ALA:HB3	1.91	0.53
1:A:32:ASN:HB3	1:B:6:ARG:HB2	1.89	0.53
1:A:46:LYS:HB3	1:A:49:ALA:HB3	1.91	0.51
1:E:19:LEU:HD22	1:E:88:ILE:HD12	1.93	0.50
1:D:122:ASP:HA	1:D:250:LYS:HD3	1.92	0.50
1:C:234:ASN:HB2	1:C:265:LEU:HD21	1.94	0.50

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:F:79:MSE:SE	1:F:130:ARG:HG2	2.63	0.49
1:A:87:PHE:HB3	1:B:89:ARG:HB3	1.95	0.48
1:B:52:LEU:HD23	1:B:99:THR:HA	1.94	0.48
1:B:234:ASN:ND2	5:B:510[A]:HOH:O	2.47	0.48
1:F:219[B]:ARG:HG3	5:F:452[B]:HOH:O	2.14	0.48
1:A:174:PRO:HD2	1:A:177:GLN:HB2	1.96	0.47
1:F:56:HIS:HE1	1:F:58:LYS:HE3	1.79	0.47
1:D:230:ALA:HB1	1:D:265:LEU:HD13	1.97	0.46
1:A:32:ASN:ND2	1:E:237:GLY:O	2.48	0.46
1:B:82:PRO:HB3	1:B:99:THR:O	2.15	0.46
1:C:23:LEU:HD22	1:C:43:LEU:HD22	1.96	0.46
1:D:196:HIS:HA	1:D:220:VAL:HG22	1.99	0.45
1:E:72:LEU:HD12	1:E:81:LEU:HD23	1.99	0.45
1:E:124:LEU:HD13	1:E:232:LEU:HD11	1.98	0.44
1:E:256:PRO:HB2	1:E:258:MSE:HE2	1.99	0.44
1:B:217:VAL:HG22	1:B:220:VAL:HG21	2.00	0.44
1:D:241:PRO:HA	1:D:244:GLN:HE21	1.83	0.44
1:A:82:PRO:HB3	1:A:99:THR:O	2.18	0.43
1:B:217:VAL:HG22	1:B:220:VAL:CG2	2.48	0.43
1:E:19:LEU:H	1:E:19:LEU:HD23	1.84	0.43
1:D:246:ARG:HE	1:D:250:LYS:NZ	2.17	0.43
1:A:89:ARG:HG2	1:B:69:MSE:HE3	2.00	0.42
1:A:79:MSE:HE2	1:A:131:LEU:HB2	2.01	0.42
1:A:258:MSE:SE	1:A:261:LEU:CD2	3.14	0.42
1:E:130:ARG:NH2	5:E:445:HOH:O	2.52	0.42
1:D:204:LEU:HB3	1:D:211:LEU:HD11	2.00	0.42
1:A:89:ARG:NH1	1:A:90:THR:O	2.48	0.42
1:D:232:LEU:HD23	1:D:247:LEU:HD11	2.02	0.42
1:A:19:LEU:HG	1:A:21:ALA:H	1.85	0.42
1:A:269:GLU:OE1	2:A:301:KAN:N3	2.51	0.42
1:C:70:VAL:HG11	1:D:70:VAL:CG1	2.50	0.42
1:F:256:PRO:HB2	1:F:258:MSE:CE	2.50	0.42
1:B:10:CYS:HB2	1:E:114:PRO:HG3	2.01	0.41
1:D:204:LEU:HD22	1:D:211:LEU:HD21	2.03	0.41
1:D:181:GLU:HB2	1:D:266:MSE:HE1	2.02	0.41
1:F:256:PRO:HB2	1:F:258:MSE:HE2	2.02	0.41
1:D:230:ALA:HA	1:D:261:LEU:HD12	2.03	0.41
1:B:43:LEU:HB2	1:B:52:LEU:HB2	2.02	0.41
1:D:69:MSE:HG3	1:D:87:PHE:CG	2.55	0.41
1:C:70:VAL:CG1	1:D:70:VAL:HG11	2.50	0.41
1:A:83[B]:THR:HB	5:A:443:HOH:O	2.20	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:46:LYS:HB3	1:B:49:ALA:HB3	2.02	0.41
1:B:106:ALA:O	1:B:110:LEU:HB2	2.21	0.41
1:D:199:PHE:CE2	1:D:232:LEU:HD22	2.56	0.40
1:A:42:ARG:HD2	1:A:51:GLU:OE2	2.21	0.40
1:D:69:MSE:HG3	1:D:87:PHE:CD1	2.56	0.40
1:B:83:THR:O	1:B:98:THR:HA	2.21	0.40
1:D:226:TYR:HA	1:D:229:LEU:HB2	2.02	0.40
1:F:148:ARG:HD2	1:F:219[A]:ARG:NH1	2.37	0.40
1:B:69:MSE:HE2	1:B:84:ILE:HD13	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	252/272 (93%)	244 (97%)	8 (3%)	0	100	100
1	B	259/272 (95%)	250 (96%)	9 (4%)	0	100	100
1	C	259/272 (95%)	249 (96%)	10 (4%)	0	100	100
1	D	238/272 (88%)	224 (94%)	14 (6%)	0	100	100
1	E	260/272 (96%)	251 (96%)	9 (4%)	0	100	100
1	F	241/272 (89%)	234 (97%)	7 (3%)	0	100	100
All	All	1509/1632 (92%)	1452 (96%)	57 (4%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	221/231 (96%)	212 (96%)	9 (4%)	37	54
1	B	233/231 (101%)	224 (96%)	9 (4%)	39	57
1	C	230/231 (100%)	224 (97%)	6 (3%)	54	73
1	D	211/231 (91%)	204 (97%)	7 (3%)	45	64
1	E	231/231 (100%)	225 (97%)	6 (3%)	54	73
1	F	216/231 (94%)	213 (99%)	3 (1%)	74	88
All	All	1342/1386 (97%)	1302 (97%)	40 (3%)	48	68

All (40) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	20	ASP
1	A	27	ARG
1	A	85	LYS
1	A	130	ARG
1	A	176	GLU
1	A	177	GLN
1	A	188	PHE
1	A	204	LEU
1	A	265	LEU
1	B	23	LEU
1	B	69	MSE
1	B	70	VAL
1	B	76	THR
1	B	110	LEU
1	B	211	LEU
1	B	217	VAL
1	B	219	ARG
1	B	262	GLN
1	C	31	ASP
1	C	110	LEU
1	C	146	VAL
1	C	204	LEU
1	C	212	ILE
1	C	238	GLU
1	D	39	THR
1	D	43	LEU
1	D	81	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	86	HIS
1	D	204	LEU
1	D	255	ASN
1	D	265	LEU
1	E	9	SER
1	E	23	LEU
1	E	33	VAL
1	E	72	LEU
1	E	116	SER
1	E	130	ARG
1	F	86	HIS
1	F	130	ARG
1	F	242	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	249	GLN
1	B	32	ASN
1	B	73	ASN
1	B	234	ASN
1	B	249	GLN
1	D	73	ASN
1	D	244	GLN
1	F	138	ASN
1	F	153	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry

15 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	KAN	A	301	-	35,35,35	0.67	0	44,52,52	1.09	4 (9%)
3	537	A	302	-	15,20,20	1.93	3 (20%)	18,30,30	2.01	2 (11%)
4	ACT	A	303	-	1,3,3	2.04	1 (100%)	0,3,3	0.00	-
2	KAN	B	301	-	35,35,35	0.67	0	44,52,52	1.34	6 (13%)
4	ACT	B	302	-	1,3,3	2.30	1 (100%)	0,3,3	0.00	-
2	KAN	C	301	-	35,35,35	0.77	1 (2%)	44,52,52	1.44	5 (11%)
3	537	C	302	-	15,20,20	1.97	3 (20%)	18,30,30	2.11	5 (27%)
4	ACT	C	303	-	1,3,3	1.31	0	0,3,3	0.00	-
4	ACT	C	304	-	1,3,3	1.28	0	0,3,3	0.00	-
2	KAN	D	301	-	35,35,35	0.71	0	44,52,52	1.10	4 (9%)
4	ACT	D	302	-	1,3,3	0.45	0	0,3,3	0.00	-
2	KAN	E	301	-	35,35,35	0.89	2 (5%)	44,52,52	1.17	7 (15%)
3	537	E	302	-	15,20,20	2.02	4 (26%)	18,30,30	2.23	3 (16%)
2	KAN	F	301	-	35,35,35	0.63	0	44,52,52	1.11	5 (11%)
4	ACT	F	302	-	1,3,3	3.56	1 (100%)	0,3,3	0.00	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	KAN	A	301	-	-	0/12/72/72	0/3/3/3
3	537	A	302	-	-	0/0/0/0	0/4/4/4
4	ACT	A	303	-	-	0/0/0/0	0/0/0/0
2	KAN	B	301	-	-	0/12/72/72	0/3/3/3
4	ACT	B	302	-	-	0/0/0/0	0/0/0/0
2	KAN	C	301	-	-	0/12/72/72	0/3/3/3
3	537	C	302	-	-	0/0/0/0	0/4/4/4

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ACT	C	303	-	-	0/0/0/0	0/0/0/0
4	ACT	C	304	-	-	0/0/0/0	0/0/0/0
2	KAN	D	301	-	-	0/12/72/72	0/3/3/3
4	ACT	D	302	-	-	0/0/0/0	0/0/0/0
2	KAN	E	301	-	-	0/12/72/72	0/3/3/3
3	537	E	302	-	-	0/0/0/0	0/4/4/4
2	KAN	F	301	-	-	0/12/72/72	0/3/3/3
4	ACT	F	302	-	-	0/0/0/0	0/0/0/0

All (16) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	302	537	N24-N23	-5.82	1.25	1.37
3	C	302	537	N24-N23	-5.80	1.25	1.37
3	A	302	537	N24-N23	-5.75	1.26	1.37
3	E	302	537	C3-C21	-2.64	1.39	1.45
3	C	302	537	C3-C21	-2.25	1.40	1.45
3	A	302	537	C3-C21	-2.15	1.40	1.45
4	A	303	ACT	CH3-C	2.04	1.51	1.48
2	E	301	KAN	O12-C13	2.06	1.47	1.41
3	E	302	537	C14-C15	2.18	1.41	1.36
3	A	302	537	C14-C15	2.19	1.41	1.36
2	C	301	KAN	O9-C1	2.25	1.47	1.41
3	E	302	537	C14-C13	2.26	1.41	1.36
3	C	302	537	C14-C15	2.28	1.41	1.36
4	B	302	ACT	CH3-C	2.30	1.52	1.48
2	E	301	KAN	O5-C1	3.26	1.50	1.41
4	F	302	ACT	CH3-C	3.56	1.53	1.48

All (41) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	301	KAN	C6-C5-C4	-4.78	104.13	113.17
2	C	301	KAN	C6-C5-C4	-4.65	104.38	113.17
3	E	302	537	C20-C16-C11	-3.28	116.65	119.14
2	D	301	KAN	C6-C5-C4	-3.12	107.28	113.17
2	D	301	KAN	C1-O9-C10	-3.06	110.01	118.01
2	F	301	KAN	C1-O9-C10	-2.92	110.39	118.01
2	B	301	KAN	O11-C13-O12	-2.77	103.68	110.68
2	B	301	KAN	C13-O11-C8	-2.50	111.49	118.01
2	E	301	KAN	O6-C2-C3	-2.39	104.95	110.34
2	E	301	KAN	C18-C17-C16	-2.34	107.25	113.02

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	301	KAN	O11-C8-C7	-2.30	103.36	108.92
2	E	301	KAN	C13-O11-C8	-2.27	112.07	118.01
3	C	302	537	C20-C16-C11	-2.26	117.43	119.14
3	C	302	537	C2-C3-C21	-2.26	119.38	122.61
2	A	301	KAN	C6-C5-C4	-2.14	109.12	113.17
2	C	301	KAN	C13-O11-C8	-2.08	112.58	118.01
2	F	301	KAN	O12-C13-C14	-2.01	106.16	110.28
2	F	301	KAN	O9-C10-C9	2.02	112.38	107.17
2	A	301	KAN	C10-C9-C8	2.02	113.14	109.06
2	F	301	KAN	O5-C5-C4	2.03	113.49	109.68
2	D	301	KAN	O5-C1-C2	2.03	114.45	110.28
2	C	301	KAN	O11-C8-C9	2.04	112.44	107.17
2	E	301	KAN	O5-C5-C4	2.16	113.74	109.68
2	D	301	KAN	C10-C9-C8	2.16	113.43	109.06
2	E	301	KAN	C1-O5-C5	2.18	117.98	113.75
2	F	301	KAN	O11-C8-C9	2.27	113.03	107.17
3	C	302	537	C2-C3-C4	2.27	121.46	118.59
2	E	301	KAN	O12-C17-C18	2.28	112.11	106.36
2	C	301	KAN	O9-C1-C2	2.30	113.70	108.10
2	A	301	KAN	O12-C17-C18	2.38	112.36	106.36
2	A	301	KAN	O9-C1-C2	2.56	114.34	108.10
2	B	301	KAN	O9-C10-C9	2.57	113.80	107.17
2	E	301	KAN	O12-C17-C16	2.66	114.68	109.68
2	B	301	KAN	O5-C5-C6	2.71	111.39	106.10
3	A	302	537	C13-C12-N24	4.36	137.56	130.22
2	C	301	KAN	O5-C5-C6	4.41	114.72	106.10
3	C	302	537	C13-C12-N24	4.43	137.69	130.22
3	E	302	537	C13-C12-N24	4.52	137.84	130.22
3	C	302	537	C15-C16-C20	5.70	125.42	119.12
3	A	302	537	C15-C16-C20	5.96	125.70	119.12
3	E	302	537	C15-C16-C20	6.59	126.40	119.12

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	KAN	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	247/272 (90%)	-0.08	6 (2%) 62 65	27, 46, 81, 95	0
1	B	257/272 (94%)	0.12	22 (8%) 13 15	30, 48, 131, 165	0
1	C	256/272 (94%)	-0.03	15 (5%) 26 30	29, 46, 108, 153	0
1	D	236/272 (86%)	0.71	37 (15%) 3 3	41, 67, 139, 197	0
1	E	258/272 (94%)	-0.11	10 (3%) 43 47	32, 46, 77, 113	0
1	F	239/272 (87%)	-0.05	9 (3%) 44 49	30, 47, 117, 148	0
All	All	1493/1632 (91%)	0.09	99 (6%) 22 25	27, 49, 112, 197	0

All (99) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	41	TYR	8.6
1	D	29	ALA	8.4
1	D	42	ARG	7.9
1	D	31	ASP	7.2
1	D	28	TRP	6.6
1	C	19	LEU	6.4
1	C	5	GLN	5.9
1	B	47	PRO	5.8
1	F	24	TYR	5.8
1	E	14	ARG	5.6
1	B	28	TRP	5.5
1	C	24	TYR	5.4
1	D	43	LEU	5.4
1	D	51	GLU	5.4
1	D	38	ALA	5.3
1	D	30	ARG	5.3
1	B	24	TYR	5.2
1	B	2	SER	5.0
1	B	36	SER	4.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	32	ASN	4.9
1	C	7	GLU	4.9
1	A	47	PRO	4.9
1	B	48	ASN	4.7
1	C	6	ARG	4.7
1	F	48	ASN	4.7
1	F	23	LEU	4.6
1	E	33	VAL	4.4
1	D	116	SER	4.3
1	B	33	VAL	4.3
1	E	36	SER	4.3
1	A	19	LEU	4.2
1	B	44	TYR	4.1
1	D	26	TYR	4.0
1	D	27	ARG	4.0
1	F	42	ARG	4.0
1	D	47	PRO	3.8
1	D	48	ASN	3.7
1	D	107	PHE	3.7
1	C	8	THR	3.7
1	D	44	TYR	3.6
1	B	4	ILE	3.5
1	C	36	SER	3.5
1	B	14	ARG	3.5
1	F	22	ASP	3.5
1	B	37	GLY	3.4
1	E	19	LEU	3.4
1	B	15	LEU	3.3
1	E	34	GLY	3.3
1	D	50	PRO	3.3
1	C	35	GLN	3.3
1	F	46	LYS	3.2
1	D	124	LEU	3.2
1	D	115	ASP	3.2
1	F	27	ARG	3.1
1	C	11	SER	3.1
1	C	34	GLY	3.1
1	D	111	GLU	3.1
1	B	3	HIS	3.1
1	D	101	ILE	3.0
1	A	172	GLY	3.0
1	B	22	ASP	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	27	ARG	2.9
1	E	35	GLN	2.9
1	D	49	ALA	2.8
1	A	48	ASN	2.8
1	D	46	LYS	2.7
1	D	119	ASN	2.7
1	E	15	LEU	2.7
1	D	249	GLN	2.7
1	B	187	PRO	2.6
1	D	255	ASN	2.6
1	B	49	ALA	2.6
1	D	187	PRO	2.6
1	D	45	GLY	2.6
1	B	25	GLY	2.6
1	B	5	GLN	2.6
1	A	173	TRP	2.5
1	D	212	ILE	2.5
1	C	27	ARG	2.5
1	C	33	VAL	2.5
1	D	254	ASP	2.5
1	C	102	PRO	2.4
1	D	232	LEU	2.4
1	D	190	PRO	2.3
1	F	44	TYR	2.3
1	D	113	TYR	2.3
1	E	113	TYR	2.2
1	B	41	TYR	2.2
1	F	26	TYR	2.2
1	E	168	ASP	2.1
1	B	50	PRO	2.1
1	A	190	PRO	2.1
1	D	256	PRO	2.1
1	D	164	SER	2.1
1	D	189	SER	2.1
1	E	18	ASN	2.1
1	D	109	VAL	2.1
1	C	47	PRO	2.0
1	C	113	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	ACT	C	303	4/4	0.93	0.15	1.41	77,77,77,77	0
2	KAN	D	301	33/33	0.93	0.18	1.20	36,57,96,300	0
4	ACT	C	304	4/4	0.96	0.17	0.50	77,78,79,79	0
2	KAN	F	301	33/33	0.96	0.10	-0.18	23,37,55,111	0
2	KAN	C	301	33/33	0.96	0.09	-0.71	21,41,63,99	0
3	537	E	302	17/17	0.99	0.08	-0.76	26,35,65,66	0
2	KAN	A	301	33/33	0.95	0.10	-0.82	32,47,71,91	0
2	KAN	E	301	33/33	0.96	0.09	-0.83	25,39,86,142	0
3	537	A	302	17/17	0.97	0.10	-1.02	32,44,83,300	0
2	KAN	B	301	33/33	0.97	0.08	-1.11	19,34,52,112	0
3	537	C	302	17/17	0.97	0.08	-1.35	30,42,75,96	0
4	ACT	D	302	4/4	0.93	0.17	-	83,83,83,83	0
4	ACT	F	302	4/4	0.91	0.16	-	74,74,74,74	0
4	ACT	A	303	4/4	0.87	0.23	-	91,91,91,92	0
4	ACT	B	302	4/4	0.93	0.08	-	77,77,78,78	0

6.5 Other polymers [i](#)

There are no such residues in this entry.