



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 09:03 AM GMT

PDB ID : 3GZN
Title : Structure of NEDD8-activating enzyme in complex with NEDD8 and MLN4924
Authors : Sintchak, M.D.
Deposited on : 2009-04-07
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	1578 (3.00-3.00)
Clashscore	102246	1912 (3.00-3.00)
Ramachandran outliers	100387	1853 (3.00-3.00)
Sidechain outliers	100360	1856 (3.00-3.00)
RSRZ outliers	91569	1592 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	534	<div> <div>83%</div> <div>15%</div> <div>••</div> </div>
1	C	534	<div> <div>86%</div> <div>12%</div> <div>••</div> </div>
2	B	463	<div> <div>77%</div> <div>15%</div> <div>• 7%</div> </div>
2	D	463	<div> <div>79%</div> <div>13%</div> <div>• 7%</div> </div>
3	I	82	<div> <div>80%</div> <div>16%</div> <div>•</div> </div>

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	J	82	 A horizontal bar chart showing the quality of chain J. The bar is divided into three segments: a green segment representing 84%, a yellow segment representing 12%, and a small grey segment at the end. A single dot is visible at the far right end of the bar.

2 Entry composition

There are 5 unique types of molecules in this entry. The entry contains 16050 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NEDD8-activating enzyme E1 regulatory subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	527	Total	C	N	O	S	0	0	0
			4095	2587	695	797	16			
1	C	526	Total	C	N	O	S	0	0	0
			4065	2573	687	789	16			

- Molecule 2 is a protein called NEDD8-activating enzyme E1 catalytic subunit.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	429	Total	C	N	O	S	0	0	0
			3311	2119	558	616	18			
2	D	429	Total	C	N	O	S	0	0	0
			3320	2127	558	617	18			

- Molecule 3 is a protein called NEDD8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
3	I	79	Total	C	N	O	S	0	0	0
			593	373	102	116	2			
3	J	79	Total	C	N	O	S	0	0	0
			602	380	103	117	2			

There are 12 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
I	-5	HIS	-	EXPRESSION TAG	UNP Q15843
I	-4	HIS	-	EXPRESSION TAG	UNP Q15843
I	-3	HIS	-	EXPRESSION TAG	UNP Q15843
I	-2	HIS	-	EXPRESSION TAG	UNP Q15843
I	-1	HIS	-	EXPRESSION TAG	UNP Q15843
I	0	HIS	-	EXPRESSION TAG	UNP Q15843
J	-5	HIS	-	EXPRESSION TAG	UNP Q15843
J	-4	HIS	-	EXPRESSION TAG	UNP Q15843

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
J	-3	HIS	-	EXPRESSION TAG	UNP Q15843
J	-2	HIS	-	EXPRESSION TAG	UNP Q15843
J	-1	HIS	-	EXPRESSION TAG	UNP Q15843
J	0	HIS	-	EXPRESSION TAG	UNP Q15843

- Molecule 4 is [(1S,2S,4R)-4-{4-[(1S)-2,3-DIHYDRO-1H-INDEN-1-YLAMINO]-7H-PYRROLO[2,3-D]PYRIMIDIN-7-YL}-2-HYDROXYCYCLOPENTYL]METHYL SULFAMATE (three-letter code: B39) (formula: C₂₁H₂₅N₅O₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
4	J	1	Total	C	N	O	S	0	0
			31	21	5	4	1		
4	I	1	Total	C	N	O	S	0	0
			31	21	5	4	1		


- Molecule 5 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Zn	0	0
			1	1		
5	D	1	Total	Zn	0	0
			1	1		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: NEDD8-activating enzyme E1 regulatory subunit




- Molecule 1: NEDD8-activating enzyme E1 regulatory subunit




- Molecule 2: NEDD8-activating enzyme E1 catalytic subunit




- Molecule 2: NEDD8-activating enzyme E1 catalytic subunit



- Molecule 3: NEDD8



- Molecule 3: NEDD8



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	135.03Å 228.72Å 229.31Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	49.36 – 3.00 49.36 – 3.00	Depositor EDS
% Data completeness (in resolution range)	97.9 (49.36-3.00) 97.3 (49.36-3.00)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.08 (at 3.01Å)	Xtriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.230 , 0.287 0.228 , 0.283	Depositor DCC
R_{free} test set	3515 reflections (5.32%)	DCC
Wilson B-factor (Å ²)	70.3	Xtriage
Anisotropy	0.085	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.34 , 30.9	EDS
Estimated twinning fraction	0.428 for -h,-l,-k	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.53$, $\langle L^2 \rangle = 0.37$	Xtriage
Outliers	2 of 69615 reflections (0.003%)	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	16050	wwPDB-VP
Average B, all atoms (Å ²)	68.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.88% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, B39

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.43	0/4177	0.55	0/5670
1	C	0.42	0/4146	0.55	0/5631
2	B	0.44	0/3388	0.59	1/4622 (0.0%)
2	D	0.43	0/3397	0.59	1/4633 (0.0%)
3	I	0.39	0/600	0.64	0/810
3	J	0.40	0/609	0.57	0/820
All	All	0.43	0/16317	0.57	2/22186 (0.0%)

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	351	LEU	CA-CB-CG	5.73	128.48	115.30
2	B	351	LEU	CA-CB-CG	5.49	127.92	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4095	0	3924	57	0
1	C	4065	0	3883	47	0
2	B	3311	0	3219	50	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
2	D	3320	0	3238	49	0
3	I	593	0	582	6	0
3	J	602	0	598	8	0
4	I	31	0	23	4	0
4	J	31	0	23	3	0
5	B	1	0	0	0	0
5	D	1	0	0	0	0
All	All	16050	0	15490	202	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 202 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:206:PHE:HB3	2:D:347:ASP:HB3	1.58	0.85
1:C:63:SER:HB3	1:C:108:SER:HB3	1.62	0.81
1:A:182:HIS:HE1	1:A:519:GLN:HE21	1.26	0.81
1:C:182:HIS:CE1	1:C:519:GLN:HE21	2.01	0.79
2:B:75:ILE:HB	2:B:164:CYS:HB3	1.65	0.78

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	525/534 (98%)	490 (93%)	32 (6%)	3 (1%)	30	72
1	C	522/534 (98%)	493 (94%)	26 (5%)	3 (1%)	30	72
2	B	425/463 (92%)	390 (92%)	30 (7%)	5 (1%)	16	56
2	D	425/463 (92%)	398 (94%)	22 (5%)	5 (1%)	16	56

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
3	I	77/82 (94%)	71 (92%)	5 (6%)	1 (1%)	15 53
3	J	77/82 (94%)	71 (92%)	5 (6%)	1 (1%)	15 53
All	All	2051/2158 (95%)	1913 (93%)	120 (6%)	18 (1%)	21 64

5 of 18 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	525	ASN
2	B	341	ASN
1	C	525	ASN
2	D	341	ASN
2	D	381	LYS

5.3.2 Protein sidechains ⓘ

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	434/464 (94%)	423 (98%)	11 (2%)	55 86
1	C	428/464 (92%)	421 (98%)	7 (2%)	70 92
2	B	357/404 (88%)	351 (98%)	6 (2%)	68 91
2	D	359/404 (89%)	354 (99%)	5 (1%)	74 93
3	I	60/72 (83%)	58 (97%)	2 (3%)	45 82
3	J	61/72 (85%)	60 (98%)	1 (2%)	70 92
All	All	1699/1880 (90%)	1667 (98%)	32 (2%)	65 90

5 of 32 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	399	SER
1	C	61	ILE
3	I	70	VAL
2	B	444	LEU
1	C	101	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 35 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	341	ASN
1	C	101	GLN
2	D	341	ASN
2	B	363	ASN
1	C	71	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	B39	I	464	3	28,35,35	3.56	6 (21%)	33,52,52	3.27	7 (21%)
4	B39	J	464	3	28,35,35	3.60	6 (21%)	33,52,52	3.33	7 (21%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	B39	I	464	3	-	0/10/35/35	0/5/5/5
4	B39	J	464	3	-	0/10/35/35	0/5/5/5

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	J	464	B39	O27-S28	-10.52	1.43	1.57
4	I	464	B39	O27-S28	-9.78	1.44	1.57
4	J	464	B39	C17-C12	-3.22	1.41	1.44
4	I	464	B39	C17-C12	-2.99	1.41	1.44
4	J	464	B39	C11-N7	-2.24	1.33	1.37

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	J	464	B39	O30-S28-O29	-13.54	107.86	119.98
4	I	464	B39	O30-S28-O29	-12.99	108.35	119.98
4	I	464	B39	N14-C15-N16	-9.83	121.37	128.89
4	J	464	B39	N14-C15-N16	-9.82	121.37	128.89
4	I	464	B39	C12-C17-N16	-3.57	118.91	121.46

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	I	464	B39	4	0
4	J	464	B39	3	0

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	527/534 (98%)	0.27	4 (0%) 87 67	42, 63, 104, 109	0
1	C	526/534 (98%)	0.28	4 (0%) 87 67	44, 63, 102, 112	0
2	B	429/463 (92%)	0.29	5 (1%) 81 55	45, 64, 100, 116	0
2	D	429/463 (92%)	0.33	8 (1%) 70 41	46, 63, 100, 118	0
3	I	79/82 (96%)	0.10	1 (1%) 79 53	47, 65, 75, 79	0
3	J	79/82 (96%)	0.13	0 100 100	48, 65, 75, 82	0
All	All	2069/2158 (95%)	0.28	22 (1%) 82 58	42, 64, 102, 118	0

The worst 5 of 22 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	294	ILE	3.2
1	C	263	ASN	2.9
2	D	281	ILE	2.7
2	B	295	ARG	2.7
2	B	294	ILE	2.6

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q<0.9
4	B39	I	464	31/31	0.96	0.28	0.28	48,56,60,60	4
4	B39	J	464	31/31	0.97	0.27	0.06	50,55,60,60	4
5	ZN	D	465	1/1	1.00	0.18	-1.02	56,56,56,56	0
5	ZN	B	465	1/1	0.98	0.19	-1.08	56,56,56,56	0

6.5 Other polymers [i](#)

There are no such residues in this entry.