



wwPDB X-ray Structure Validation Summary Report ⓘ

Feb 1, 2016 – 02:50 AM GMT

PDB ID : 2IW6
Title : STRUCTURE OF HUMAN THR160-PHOSPHO CDK2-CYCLIN A COM-
PLEXED WITH A BISANILINOPYRIMIDINE INHIBITOR
Authors : Pratt, D.J.; Bentley, J.; Jewsbury, P.; Boyle, F.T.; Endicott, J.A.; Noble,
M.E.M.
Deposited on : 2006-06-26
Resolution : 2.30 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	3852 (2.30-2.30)
Clashscore	102246	4452 (2.30-2.30)
Ramachandran outliers	100387	4410 (2.30-2.30)
Sidechain outliers	100360	4409 (2.30-2.30)
RSRZ outliers	91569	3857 (2.30-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	302	<div> <div>4%</div> <div>70%</div> <div>23%</div> <div>• •</div> </div>
1	C	302	<div> <div>20%</div> <div>74%</div> <div>19%</div> <div>5% •</div> </div>
2	B	260	<div> <div>3%</div> <div>82%</div> <div>15%</div> <div>• •</div> </div>
2	D	260	<div> <div>21%</div> <div>73%</div> <div>19%</div> <div>6% •</div> </div>

2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 9062 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called CELL DIVISION PROTEIN KINASE 2.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	293	Total	C	N	O	P	S	0	0	0
			2354	1529	399	417	1	8			
1	C	296	Total	C	N	O	P	S	0	0	0
			2378	1542	402	425	1	8			

There are 10 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	EXPRESSION TAG	UNP P24941
A	-2	PRO	-	EXPRESSION TAG	UNP P24941
A	-1	GLY	-	EXPRESSION TAG	UNP P24941
A	0	SER	-	EXPRESSION TAG	UNP P24941
A	89	THR	LYS	ENGINEERED MUTATION	UNP P24941
C	-3	GLY	-	EXPRESSION TAG	UNP P24941
C	-2	PRO	-	EXPRESSION TAG	UNP P24941
C	-1	GLY	-	EXPRESSION TAG	UNP P24941
C	0	SER	-	EXPRESSION TAG	UNP P24941
C	89	THR	LYS	ENGINEERED MUTATION	UNP P24941

- Molecule 2 is a protein called CYCLIN-A2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
2	B	257	Total	C	N	O	S	0	0	0
			2076	1345	338	382	11			
2	D	255	Total	C	N	O	S	0	0	0
			2061	1336	336	378	11			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	173	MET	-	EXPRESSION TAG	UNP P20248
D	173	MET	-	EXPRESSION TAG	UNP P20248

-
- Chemical structure of compound QQ2. The molecule features a pyridine ring (N6, C1-C5) connected via an amine group (N19, C20) to a benzene ring (C21-C26). This benzene ring is further connected to a pyrazole ring (N15, N18, C3-C4). The pyrazole ring is substituted with a chlorine atom (Cl1) and a methyl group (C16). A hydroxyl group (HO, O28) is attached to the pyrazole ring via a methylene group (C27). The structure is labeled with atom IDs: C31, N30, C32, C33, O28, C29, C27, O26, C23, C22, C21, C20, C25, C24, N19, N6, C1, N2, C4, C3, N7, C16, C17, N18, Cl1, C13, C12, C11, C10, C9, C8.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total 33	C 24	Cl 1	N 6	O 2	0	0
3	C	1	Total 33	C 24	Cl 1	N 6	O 2	0	0

- SGM
-
- The diagram shows the chemical structure of (S)-butane-2-thiol. The carbon chain is represented by a zigzag line. The chiral center is labeled C2(R) in green. The atoms are color-coded: carbons are green, hydrogens are red, and the sulfur atom is yellow. The groups are labeled as follows: C3 is the carbon to the left of C2, C1 is the carbon to the right of C2, and S1 is the sulfur atom bonded to C1. The hydrogens are labeled H03 and H02 in red. The structure is drawn in a perspective view, with the C2-C3 bond and the C2-H03 bond in the plane of the paper, the C2-H02 bond coming out of the plane (indicated by a wedge), and the C2-S1 bond going into the plane (indicated by a dashed line).
- CC(C)CS

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
4	B	1	Total	C	O	S	0	0
			6	3	2	1		
4	D	1	Total	C	O	S	0	0
			6	3	2	1		

- Molecule 5 is MAGNESIUM ION (three-letter code: MG) (formula: Mg).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	Mg	0	0
			1	1		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	48	Total	O	0	0
			48	48		
6	B	49	Total	O	0	0
			49	49		
6	C	9	Total	O	0	0
			9	9		
6	D	8	Total	O	0	0
			8	8		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: CELL DIVISION PROTEIN KINASE 2




• Molecule 1: CELL DIVISION PROTEIN KINASE 2



• Molecule 2: CYCLIN-A2



• Molecule 2: CYCLIN-A2



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	73.78Å 134.56Å 148.26Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	100.00 – 2.30 41.06 – 2.30	Depositor EDS
% Data completeness (in resolution range)	93.1 (100.00-2.30) 92.7 (41.06-2.30)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.07 (at 2.29Å)	Xtriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.228 , 0.287 0.218 , 0.273	Depositor DCC
R_{free} test set	3125 reflections (5.37%)	DCC
Wilson B-factor (Å ²)	41.4	Xtriage
Anisotropy	0.425	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 49.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	1 of 61338 reflections (0.002%)	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	9062	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.97% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: TPO, MG, SGM, QQ2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.83	0/2403	0.88	2/3260 (0.1%)
1	C	0.63	0/2428	0.74	2/3296 (0.1%)
2	B	0.79	1/2126 (0.0%)	0.83	0/2886
2	D	1.23	26/2110 (1.2%)	0.82	2/2864 (0.1%)
All	All	0.89	27/9067 (0.3%)	0.82	6/12306 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	D	0	1

The worst 5 of 27 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	375	SER	CB-OG	19.35	1.67	1.42
2	D	423	LEU	C-O	13.67	1.49	1.23
2	D	370	GLN	CD-OE1	10.37	1.46	1.24
2	D	429	THR	C-N	9.75	1.56	1.34
2	D	374	GLU	CD-OE2	9.52	1.36	1.25

The worst 5 of 6 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	76	LEU	CA-CB-CG	5.83	128.71	115.30
2	D	423	LEU	O-C-N	5.74	131.89	122.70
1	A	230	VAL	CB-CA-C	-5.67	100.63	111.40
1	A	78	LEU	CA-CB-CG	5.33	127.56	115.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	429	THR	CA-C-N	-5.31	105.52	117.20

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	D	428	GLU	Mainchain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2354	0	2395	57	0
1	C	2378	0	2413	41	0
2	B	2076	0	2098	36	0
2	D	2061	0	2086	38	0
3	A	33	0	27	4	0
3	C	33	0	27	4	0
4	B	6	0	7	0	0
4	D	6	0	7	1	0
5	B	1	0	0	0	0
6	A	48	0	0	3	0
6	B	49	0	0	1	0
6	C	9	0	0	1	0
6	D	8	0	0	1	0
All	All	9062	0	9060	162	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

The worst 5 of 162 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:414:LYS:CE	2:D:414:LYS:NZ	1.70	1.53
2:D:375:SER:OG	2:D:375:SER:CB	1.67	1.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:193:CYS:SG	4:D:1433:SGM:S1	2.30	1.29
2:D:361:HIS:HD2	2:D:391:LEU:HD21	1.20	1.07
1:A:284:PRO:O	1:A:287:GLN:HG2	1.59	1.03

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	288/302 (95%)	277 (96%)	10 (4%)	1 (0%)	46	57
1	C	293/302 (97%)	268 (92%)	22 (8%)	3 (1%)	19	21
2	B	255/260 (98%)	250 (98%)	5 (2%)	0	100	100
2	D	253/260 (97%)	242 (96%)	10 (4%)	1 (0%)	39	48
All	All	1089/1124 (97%)	1037 (95%)	47 (4%)	5 (0%)	34	41

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	C	164	VAL
1	C	167	TRP
1	A	164	VAL
1	C	40	GLU
2	D	372	TRP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	257/264 (97%)	230 (90%)	27 (10%)	8	9
1	C	260/264 (98%)	230 (88%)	30 (12%)	7	7
2	B	231/234 (99%)	221 (96%)	10 (4%)	35	47
2	D	229/234 (98%)	215 (94%)	14 (6%)	23	30
All	All	977/996 (98%)	896 (92%)	81 (8%)	14	17

5 of 81 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
2	B	389	PRO
1	C	76	LEU
2	D	348	LEU
1	C	0	SER
1	C	46	SER

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 25 such sidechains are listed below:

Mol	Chain	Res	Type
2	B	425	ASN
1	C	60	HIS
2	D	361	HIS
1	C	59	ASN
1	C	84	HIS

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected

value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
1	TPO	A	160	1	8,10,11	0.88	0	7,14,16	1.21	1 (14%)
1	TPO	C	160	1	8,10,11	0.73	0	7,14,16	1.27	1 (14%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	TPO	A	160	1	-	0/8/11/13	0/0/0/0
1	TPO	C	160	1	-	0/8/11/13	0/0/0/0

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	C	160	TPO	O-C-CA	-2.25	119.48	125.44
1	A	160	TPO	O3P-P-O2P	2.41	116.56	107.38

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 5 ligands modelled in this entry, 1 is monoatomic - leaving 4 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length

(or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
3	QQ2	A	1296	-	34,35,35	1.05	2 (5%)	42,47,47	2.39	14 (33%)
4	SGM	B	1433	2	5,5,5	0.99	0	5,5,5	0.94	0
3	QQ2	C	1296	-	34,35,35	0.91	3 (8%)	42,47,47	2.14	9 (21%)
4	SGM	D	1433	-	5,5,5	0.45	0	5,5,5	0.74	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	QQ2	A	1296	-	-	0/23/24/24	0/3/3/3
4	SGM	B	1433	2	-	0/4/4/4	0/0/0/0
3	QQ2	C	1296	-	-	0/23/24/24	0/3/3/3
4	SGM	D	1433	-	-	0/4/4/4	0/0/0/0

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	C	1296	QQ2	C20-N19	-2.42	1.35	1.40
3	C	1296	QQ2	C3-N7	-2.02	1.36	1.40
3	A	1296	QQ2	C5-N19	2.18	1.42	1.38
3	A	1296	QQ2	C16-C17	2.35	1.50	1.47
3	C	1296	QQ2	C13-CL1	2.50	1.79	1.73

The worst 5 of 23 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	1296	QQ2	C4-C5-N6	-4.60	117.91	123.06
3	C	1296	QQ2	N6-C1-N2	-4.46	121.16	128.67
3	C	1296	QQ2	C16-C17-N18	-3.55	172.83	177.99
3	A	1296	QQ2	N6-C1-N2	-3.44	122.87	128.67
3	A	1296	QQ2	C4-C3-N2	-3.25	116.68	122.49

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

3 monomers are involved in 9 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1296	QQ2	4	0
3	C	1296	QQ2	4	0
4	D	1433	SGM	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ > 2	OWAB(Å ²)	Q < 0.9
1	A	292/302 (96%)	0.49	13 (4%) 37 46	26, 35, 46, 59	0
1	C	295/302 (97%)	1.18	60 (20%) 1 2	24, 35, 45, 58	0
2	B	257/260 (98%)	0.31	9 (3%) 48 56	29, 35, 44, 52	0
2	D	255/260 (98%)	1.07	54 (21%) 1 2	31, 36, 43, 48	0
All	All	1099/1124 (97%)	0.77	136 (12%) 5 8	24, 35, 45, 59	0

The worst 5 of 136 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	D	367	VAL	10.9
1	C	14	THR	9.7
1	C	234	PRO	7.8
2	D	368	THR	7.7
1	C	15	TYR	7.7

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q < 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å ²)	Q < 0.9
1	TPO	A	160	11/12	0.99	0.13	-	29,32,34,34	0
1	TPO	C	160	11/12	0.97	0.11	-	31,33,36,36	0

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(\AA^2)	Q<0.9
3	QQ2	C	1296	33/33	0.85	0.19	0.11	40,44,54,56	0
4	SGM	D	1433	6/6	0.92	0.14	-0.16	54,55,55,56	0
3	QQ2	A	1296	33/33	0.91	0.16	-0.17	35,45,54,56	0
5	MG	B	1434	1/1	0.94	0.08	-1.74	31,31,31,31	0
4	SGM	B	1433	6/6	0.95	0.10	-1.82	42,45,48,50	0

6.5 Other polymers [i](#)

There are no such residues in this entry.