



# Full wwPDB X-ray Structure Validation Report ⓘ

Aug 29, 2016 – 11:26 AM EDT

PDB ID : 5IW4  
Title : Crystal structure of E. coli NudC in complex with NAD  
Authors : Li, S.; Du, J.; Patel, D.J.  
Deposited on : 2016-03-22  
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<http://wwpdb.org/validation/2016/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : unknown  
Xtriage (Phenix) : 1.9-1692  
EDS : rb-20027939  
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)  
Refmac : 5.8.0135  
CCP4 : 6.5.0  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20027939




# 1 Overall quality at a glance ⓘ

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	91344	2328 (2.60-2.60)
Clashscore	102246	2679 (2.60-2.60)
Ramachandran outliers	100387	2635 (2.60-2.60)
Sidechain outliers	100360	2635 (2.60-2.60)
RSRZ outliers	91569	2334 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	258	<div> <div>2%</div> <div>86%</div> <div>12%</div> <div>.</div> </div>
1	B	258	<div> <div>2%</div> <div>78%</div> <div>19%</div> <div>.</div> </div>

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:



Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	NAD	B	501	-	-	-	X



## 2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 4407 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NADH pyrophosphatase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	A	257	Total	C	N	O	S	0	0	0
			2086	1328	367	377	14			
1	B	258	Total	C	N	O	S	0	0	0
			2095	1333	368	380	14			

There are 2 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	0	SER	-	expression tag	UNP D6JHV3
B	0	SER	-	expression tag	UNP D6JHV3

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total	Zn	0	0
			1	1		
2	A	1	Total	Zn	0	0
			1	1		

- Molecule 3 is NICOTINAMIDE-ADENINE-DINUCLEOTIDE (three-letter code: NAD) (formula: C<sub>21</sub>H<sub>27</sub>N<sub>7</sub>O<sub>14</sub>P<sub>2</sub>).





Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	A	1	Total	C	N	O	P	0	0
			44	21	7	14	2		
3	B	1	Total	C	N	O	P	0	0
			44	21	7	14	2		

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	76	Total	O	0	0
			76	76		
4	B	60	Total	O	0	0
			60	60		




### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: NADH pyrophosphatase



- Molecule 1: NADH pyrophosphatase





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	51.93Å 99.37Å 113.66Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	45.53 – 2.60 45.52 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.7 (45.53-2.60) 99.7 (45.52-2.60)	Depositor EDS
$R_{merge}$	0.11	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.73 (at 2.61Å)	Xtriage
Refinement program	PHENIX 1.8.2_1309	Depositor
R, $R_{free}$	0.200 , 0.248 0.195 , 0.248	Depositor DCC
$R_{free}$ test set	957 reflections (5.13%)	DCC
Wilson B-factor (Å <sup>2</sup> )	43.8	Xtriage
Anisotropy	0.223	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 35.2	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	4407	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	49.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.22% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ZN, NAD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z  > 5$	RMSZ	$\# Z  > 5$
1	A	0.58	0/2142	0.50	0/2909
1	B	0.53	0/2151	0.56	1/2921 (0.0%)
All	All	0.56	0/4293	0.53	1/5830 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	217	PRO	N-CD-CG	-6.05	94.13	103.20

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2086	0	2031	21	0
1	B	2095	0	2037	44	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
3	A	44	0	26	3	0
3	B	44	0	26	4	0
4	A	76	0	0	0	0
4	B	60	0	0	1	0

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
All	All	4407	0	4120	67	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (67) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:237:PRO:HG3	1:B:242:ARG:NH2	1.50	1.24
1:B:3:ARG:NH2	1:B:9:ASP:OD1	1.77	1.18
1:B:237:PRO:CG	1:B:242:ARG:NH2	2.19	1.05
1:B:237:PRO:HG3	1:B:242:ARG:HH21	0.96	1.00
1:B:237:PRO:CG	1:B:242:ARG:HH21	1.82	0.89
1:B:3:ARG:HH21	1:B:9:ASP:CG	1.77	0.86
1:B:237:PRO:CD	1:B:242:ARG:NH2	2.44	0.80
1:A:95:HIS:HB3	1:A:105:MET:HE3	1.65	0.79
1:B:237:PRO:HD3	1:B:242:ARG:NH2	1.98	0.77
1:B:239:THR:HB	3:B:501:NAD:H72N	1.51	0.74
1:B:3:ARG:NH2	1:B:9:ASP:CG	2.39	0.74
1:B:8:LEU:C	1:B:8:LEU:HD12	2.11	0.72
1:B:59:GLN:OE1	1:B:61:ARG:NH1	2.24	0.69
1:B:237:PRO:CG	1:B:242:ARG:HH22	2.05	0.69
1:A:151:ASN:N	1:A:151:ASN:OD1	2.26	0.66
1:A:2:ASP:OD2	1:A:42:ARG:NH1	2.30	0.65
1:B:8:LEU:C	1:B:8:LEU:CD1	2.70	0.61
1:B:243:ARG:NE	1:B:247:ASP:OD2	2.26	0.59
1:B:151:ASN:OD1	1:B:151:ASN:N	2.35	0.58
1:B:74:LEU:HD13	1:B:78:LEU:HD23	1.85	0.58
1:B:22:TRP:CE2	1:B:24:PRO:HG3	2.39	0.58
1:B:122:ARG:NH2	1:B:124:TYR:OH	2.39	0.56
1:B:141:ILE:HD12	1:B:248:THR:HG21	1.88	0.56
1:B:8:LEU:O	1:B:8:LEU:HD12	2.06	0.55
1:A:239:THR:HB	3:A:501:NAD:H72N	1.72	0.55
1:A:171:VAL:HG11	1:A:186:LEU:HD21	1.88	0.54
1:B:209:ASP:OD1	1:B:209:ASP:C	2.45	0.54
3:A:501:NAD:O2N	3:A:501:NAD:H51A	2.08	0.54
1:B:80:GLN:NE2	1:B:192:GLN:OE1	2.34	0.53
1:A:228:TYR:HE1	1:A:253:ARG:HD2	1.74	0.53
1:A:148:ARG:NH2	1:A:219:GLU:OE2	2.43	0.52
1:B:228:TYR:HB2	1:B:249:VAL:HG22	1.93	0.51
1:A:13:TRP:HB2	1:A:29:PRO:HB2	1.93	0.51

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:239:THR:HB	3:B:501:NAD:N7N	2.23	0.51
1:B:197:PRO:HG2	1:B:198:GLN:HG2	1.92	0.51
1:B:135:ILE:HD12	1:B:142:LEU:HD23	1.92	0.51
1:A:231:LEU:HD11	1:A:249:VAL:HG21	1.95	0.48
1:B:195:PRO:HD2	1:B:196:PHE:CE2	2.48	0.48
1:B:143:LEU:HB3	1:B:154:HIS:HB3	1.95	0.47
1:B:194:TRP:CD1	3:B:501:NAD:H1D	2.50	0.47
1:A:136:ARG:NH2	1:A:255:GLU:OE1	2.48	0.47
1:A:239:THR:HB	3:A:501:NAD:N7N	2.30	0.47
1:B:231:LEU:HD11	1:B:249:VAL:HG21	1.95	0.47
1:B:71:VAL:HG23	1:B:71:VAL:O	2.15	0.46
1:A:131:ILE:HD13	1:A:171:VAL:HA	1.97	0.45
1:B:71:VAL:CG2	1:B:71:VAL:O	2.64	0.45
1:B:14:VAL:HG11	1:B:85:GLY:HA3	1.99	0.45
1:A:17:HIS:HB3	1:A:60:GLN:HB3	2.00	0.44
1:B:187:ARG:NH1	4:B:611:HOH:O	2.49	0.44
1:A:126:GLN:O	1:A:198:GLN:NE2	2.49	0.44
1:B:174:GLU:HG3	1:B:178:GLU:OE2	2.17	0.44
1:A:10:HIS:O	1:A:53:PRO:HG2	2.17	0.44
1:B:178:GLU:HG2	1:B:219:GLU:HG2	2.00	0.44
1:A:54:VAL:HG21	1:A:81:LEU:HD13	2.00	0.43
1:B:136:ARG:NH1	1:B:251:MET:HG2	2.34	0.43
1:B:10:HIS:ND1	1:B:52:GLU:OE1	2.43	0.42
1:B:68:VAL:O	1:B:71:VAL:HG13	2.19	0.42
1:B:22:TRP:CD2	1:B:61:ARG:HG3	2.54	0.42
1:A:148:ARG:HH21	1:A:148:ARG:HG2	1.85	0.42
1:B:244:LEU:HA	1:B:244:LEU:HD23	1.93	0.42
1:A:148:ARG:HG2	1:A:148:ARG:NH2	2.35	0.41
1:B:131:ILE:HD13	1:B:171:VAL:HA	2.02	0.41
1:B:78:LEU:HD12	1:B:78:LEU:HA	1.94	0.41
1:A:155:THR:OG1	1:A:156:VAL:N	2.54	0.41
1:A:192:GLN:HA	1:A:193:PRO:HD3	1.96	0.40
3:B:501:NAD:H8A	3:B:501:NAD:H2B	1.81	0.40
1:A:2:ASP:CG	1:A:42:ARG:HH11	2.24	0.40

There are no symmetry-related clashes.



## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	255/258 (99%)	248 (97%)	6 (2%)	1 (0%)	39	65
1	B	256/258 (99%)	248 (97%)	8 (3%)	0	100	100
All	All	511/516 (99%)	496 (97%)	14 (3%)	1 (0%)	52	77

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	232	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	221/222 (100%)	215 (97%)	6 (3%)	52	79
1	B	222/222 (100%)	210 (95%)	12 (5%)	27	52
All	All	443/444 (100%)	425 (96%)	18 (4%)	37	66

All (18) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	42	ARG
1	A	136	ARG
1	A	148	ARG
1	A	218	LYS
1	A	222	GLU

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type
1	A	243	ARG
1	B	1	MET
1	B	3	ARG
1	B	8	LEU
1	B	37	ASP
1	B	39	VAL
1	B	71	VAL
1	B	147	THR
1	B	151	ASN
1	B	182	LYS
1	B	242	ARG
1	B	251	MET
1	B	253	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 2 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).



Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	NAD	A	501	-	42,48,48	3.36	18 (42%)	46,73,73	3.10	16 (34%)
3	NAD	B	501	-	42,48,48	3.28	15 (35%)	46,73,73	2.57	12 (26%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAD	A	501	-	-	0/22/62/62	0/5/5/5
3	NAD	B	501	-	-	0/22/62/62	0/5/5/5

All (33) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	501	NAD	C4N-C3N	-10.68	1.21	1.39
3	B	501	NAD	C4N-C3N	-9.60	1.23	1.39
3	A	501	NAD	C5N-C4N	-6.32	1.26	1.38
3	A	501	NAD	C2D-C1D	-5.52	1.44	1.53
3	A	501	NAD	C2B-C1B	-5.33	1.45	1.53
3	B	501	NAD	C5N-C4N	-5.04	1.29	1.38
3	B	501	NAD	C2D-C1D	-4.44	1.46	1.53
3	A	501	NAD	C2N-C3N	-4.39	1.32	1.39
3	A	501	NAD	C2D-C3D	-4.25	1.42	1.53
3	A	501	NAD	C2B-C3B	-4.03	1.42	1.53
3	B	501	NAD	C2B-C1B	-3.81	1.47	1.53
3	B	501	NAD	C2N-C3N	-3.30	1.33	1.39
3	A	501	NAD	C5A-N7A	-2.83	1.29	1.39
3	B	501	NAD	C2D-C3D	-2.78	1.45	1.53
3	A	501	NAD	C3B-C4B	-2.77	1.45	1.53
3	A	501	NAD	C8A-N7A	-2.72	1.29	1.34
3	B	501	NAD	C2B-C3B	-2.31	1.47	1.53
3	A	501	NAD	C3D-C4D	-2.22	1.47	1.53
3	A	501	NAD	O7N-C7N	-2.22	1.19	1.24
3	B	501	NAD	O4D-C4D	2.44	1.50	1.45
3	B	501	NAD	C2A-N3A	2.72	1.37	1.32
3	B	501	NAD	C6N-C5N	2.88	1.44	1.38
3	A	501	NAD	C2A-N3A	3.51	1.38	1.32
3	A	501	NAD	O4B-C1B	4.74	1.48	1.41
3	A	501	NAD	C2N-N1N	4.93	1.42	1.35
3	A	501	NAD	C7N-N7N	5.33	1.43	1.33
3	A	501	NAD	C6N-N1N	5.53	1.51	1.35

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	501	NAD	O4D-C1D	6.10	1.49	1.41
3	B	501	NAD	C7N-N7N	6.30	1.45	1.33
3	B	501	NAD	C6N-N1N	6.46	1.53	1.35
3	B	501	NAD	C2N-N1N	6.73	1.44	1.35
3	B	501	NAD	O4B-C1B	6.91	1.51	1.41
3	B	501	NAD	O4D-C1D	7.02	1.51	1.41

All (28) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	501	NAD	N3A-C2A-N1A	-15.28	116.87	128.87
3	B	501	NAD	N3A-C2A-N1A	-11.04	120.20	128.87
3	B	501	NAD	C4D-O4D-C1D	-7.93	101.23	109.64
3	A	501	NAD	C4D-O4D-C1D	-5.47	103.85	109.64
3	A	501	NAD	O2D-C2D-C3D	-4.13	98.51	111.86
3	A	501	NAD	O3D-C3D-C2D	-3.82	99.51	111.86
3	A	501	NAD	C4B-O4B-C1B	-3.56	105.87	109.64
3	A	501	NAD	C5N-C6N-N1N	-3.40	114.58	120.46
3	B	501	NAD	C5N-C6N-N1N	-3.33	114.70	120.46
3	B	501	NAD	C4B-O4B-C1B	-3.13	106.32	109.64
3	A	501	NAD	O2B-C2B-C3B	-3.13	101.73	111.86
3	A	501	NAD	O3B-C3B-C2B	-2.80	102.81	111.86
3	A	501	NAD	O7N-C7N-N7N	-2.45	119.09	122.58
3	A	501	NAD	C2D-C1D-N1N	-2.41	108.81	113.53
3	B	501	NAD	O7N-C7N-N7N	-2.00	119.72	122.58
3	B	501	NAD	C2D-C3D-C4D	2.01	106.74	102.64
3	A	501	NAD	C3N-C7N-N7N	2.07	120.17	117.82
3	B	501	NAD	C5N-C4N-C3N	2.13	122.89	120.35
3	B	501	NAD	C3N-C2N-N1N	2.20	122.85	120.34
3	A	501	NAD	C6N-C5N-C4N	2.28	122.90	119.43
3	B	501	NAD	C2B-C3B-C4B	2.29	107.32	102.64
3	A	501	NAD	C2A-N1A-C6A	2.45	123.14	118.77
3	A	501	NAD	C5N-C4N-C3N	2.56	123.40	120.35
3	B	501	NAD	C6N-C5N-C4N	2.69	123.52	119.43
3	A	501	NAD	O4B-C1B-N9A	2.95	113.68	108.11
3	B	501	NAD	O4D-C1D-N1N	3.99	112.41	108.10
3	B	501	NAD	C2N-C3N-C4N	5.08	124.03	118.27
3	A	501	NAD	C2N-C3N-C4N	5.56	124.58	118.27

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.



2 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	501	NAD	3	0
3	B	501	NAD	4	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	257/258 (99%)	-0.18	4 (1%) 74 69	28, 45, 72, 99	0
1	B	258/258 (100%)	-0.15	4 (1%) 74 69	30, 47, 78, 105	0
All	All	515/516 (99%)	-0.17	8 (1%) 74 69	28, 46, 75, 105	0

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	120	ARG	4.7
1	B	149	HIS	4.0
1	B	7	LYS	4.0
1	A	120	ARG	3.7
1	A	228	TYR	2.7
1	A	149	HIS	2.6
1	A	59	GLN	2.4
1	B	62	ARG	2.4

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. LLDF column lists the quality of electron density of the group with respect to its neighbouring residues in protein, DNA or RNA chains.



The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	LLDF	B-factors(Å <sup>2</sup> )	Q<0.9
3	NAD	B	501	44/44	0.73	0.43	4.12	30,49,106,108	44
3	NAD	A	501	44/44	0.89	0.18	0.02	35,61,95,149	0
2	ZN	A	500	1/1	0.98	0.07	-1.67	56,56,56,56	0
2	ZN	B	500	1/1	0.98	0.09	-2.32	49,49,49,49	0

## 6.5 Other polymers ⓘ

There are no such residues in this entry.