



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 1, 2016 – 06:38 AM GMT

PDB ID : 2XUB
Title : HUMAN RPC62 SUBUNIT STRUCTURE
Authors : Lefevre, S.; Legrand, P.; Fribourg, S.
Deposited on : 2010-10-18
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.
We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
<http://wwpdb.org/validation/2016/XrayValidationReportHelp>
with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7 (RC4), CSD as536be (2015)
Xtriage (Phenix) : 1.9-1692
EDS : rb-20026688
Percentile statistics : 20151230.v01 (using entries in the PDB archive December 30th 2015)
Refmac : 5.8.0135
CCP4 : 6.5.0
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk26865


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	91344	2393 (2.80-2.80)
Clashscore	102246	2827 (2.80-2.80)
Ramachandran outliers	100387	2782 (2.80-2.80)
Sidechain outliers	100360	2784 (2.80-2.80)
RSRZ outliers	91569	2404 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	534	<div> <div>11%</div> <div>73%</div> <div>7%</div> <div>19%</div> </div>

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 6925 atoms, of which 3476 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DNA-DIRECTED RNA POLYMERASE III SUBUNIT RPC3.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace
1	A	433	Total	C	H	N	O	S	0	0	0
			6912	2162	3476	597	654	23			


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	13	Total	O	0	0
			13	13		

3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of errors displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($\text{RSRZ} > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DNA-DIRECTED RNA POLYMERASE III SUBUNIT RPC3



4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, α , β , γ	161.79Å 69.38Å 75.40Å 90.00° 113.20° 90.00°	Depositor
Resolution (Å)	43.13 – 2.80 43.13 – 2.80	Depositor EDS
% Data completeness (in resolution range)	99.5 (43.13-2.80) 99.5 (43.13-2.80)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.50 (at 2.81Å)	Xtriage
Refinement program	BUSTER 2.9.1	Depositor
R, R_{free}	0.188 , 0.230 0.204 , 0.245	Depositor DCC
R_{free} test set	943 reflections (5.22%)	DCC
Wilson B-factor (Å ²)	67.1	Xtriage
Anisotropy	0.662	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.40 , 77.5	EDS
Estimated twinning fraction	No twinning to report.	Xtriage
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtriage
Outliers	1 of 19023 reflections (0.005%)	Xtriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	6925	wwPDB-VP
Average B, all atoms (Å ²)	101.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.90% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.375 respectively for untwinned datasets, and 0.333, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\# Z > 5$	RMSZ	$\# Z > 5$
1	A	0.51	0/3480	0.68	0/4690

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3436	3476	3474	11	0
2	A	13	0	0	0	0
All	All	3449	3476	3474	11	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (11) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:278:THR:HG21	1:A:285:THR:HA	1.89	0.54
1:A:158:GLN:HG2	1:A:238:ASN:HB2	1.89	0.52
1:A:14:GLN:HG2	1:A:22:GLU:HG3	1.96	0.48
1:A:328:ASP:HA	1:A:332:GLY:HA2	1.95	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:31:THR:OG1	1:A:34:GLN:NE2	2.44	0.45
1:A:9:CYS:HG	1:A:79:CYS:HG	1.63	0.45
1:A:67:VAL:HG22	1:A:73:VAL:HG22	1.98	0.44
1:A:377:GLN:HA	1:A:380:VAL:HB	2.01	0.43
1:A:23:LYS:HD3	1:A:43:THR:HB	1.99	0.42
1:A:330:GLY:HA2	1:A:331:GLY:HA2	1.66	0.42
1:A:428:ILE:HG13	1:A:428:ILE:H	1.63	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	423 / 534 (79%)	406 (96%)	13 (3%)	4 (1%)	21 55

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	69	LYS
1	A	135	ASP
1	A	138	THR
1	A	428	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	382/476 (80%)	362 (95%)	20 (5%)	29 62

All (20) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1	MET
1	A	36	LEU
1	A	84	ARG
1	A	119	SER
1	A	134	GLU
1	A	138	THR
1	A	143	GLU
1	A	157	VAL
1	A	231	ASP
1	A	253	SER
1	A	260	ASP
1	A	291	ASN
1	A	318	ASP
1	A	325	LYS
1	A	365	ILE
1	A	368	LEU
1	A	402	ASN
1	A	428	ILE
1	A	484	LEU
1	A	502	LEU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA ⓘ

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates ⓘ

There are no carbohydrates in this entry.

5.6 Ligand geometry

There are no ligands in this entry.

5.7 Other polymers

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data ⓘ

6.1 Protein, DNA and RNA chains ⓘ

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	433/534 (81%)	1.07	60 (13%) 4 2	58, 94, 168, 209	0

All (60) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	368	LEU	5.6
1	A	386	ILE	5.2
1	A	383	PHE	4.9
1	A	391	ALA	4.9
1	A	385	MET	4.9
1	A	380	VAL	4.8
1	A	378	LYS	4.6
1	A	396	TYR	4.6
1	A	382	ASP	4.5
1	A	426	VAL	4.3
1	A	389	LYS	4.1
1	A	308	LEU	3.8
1	A	392	LYS	3.5
1	A	135	ASP	3.4
1	A	471	ILE	3.2
1	A	312	LEU	3.2
1	A	334	TYR	3.1
1	A	470	ILE	3.1
1	A	338	LEU	3.1
1	A	370	LEU	3.0
1	A	388	ALA	3.0
1	A	6	ILE	2.9
1	A	390	GLU	2.9
1	A	398	MET	2.9
1	A	397	LYS	2.8
1	A	315	LEU	2.8
1	A	379	GLN	2.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	403	PHE	2.7
1	A	269	THR	2.6
1	A	358	PHE	2.6
1	A	266	ILE	2.5
1	A	126	ALA	2.4
1	A	377	GLN	2.4
1	A	297	LEU	2.4
1	A	526	TYR	2.4
1	A	107	VAL	2.4
1	A	393	ASP	2.3
1	A	336	ILE	2.3
1	A	502	LEU	2.3
1	A	340	LYS	2.2
1	A	35	PRO	2.2
1	A	121	VAL	2.2
1	A	323	VAL	2.2
1	A	428	ILE	2.2
1	A	111	LEU	2.2
1	A	150	ARG	2.2
1	A	458	LYS	2.2
1	A	301	TYR	2.1
1	A	435	LEU	2.1
1	A	285	THR	2.1
1	A	310	GLN	2.1
1	A	314	LEU	2.1
1	A	487	ILE	2.0
1	A	151	LEU	2.0
1	A	484	LEU	2.0
1	A	376	GLU	2.0
1	A	68	HIS	2.0
1	A	155	HIS	2.0
1	A	530	THR	2.0
1	A	77	ALA	2.0

6.2 Non-standard residues in protein, DNA, RNA chains ⓘ

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates ⓘ

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.